

GRAPEVINE NAMED 'SUGRATWENTYFOUR'

5 LATIN NAME OF THE GENUS AND SPECIES CLAIMED

Vitis vinifera

VARIETY DENOMINATION

10

'SUGRATWENTYFOUR'

BACKGROUND AND SUMMARY OF THE INVENTION

15 This invention relates to the discovery and asexual propagation of a new variety of grapevine as herein described and illustrated. The new variety is characterized by medium-sized, white seedless berries having a desirable distinctive muscat flavor, and a crisp juicy flesh. The grapes are round with a tough skin that is greenish yellow in color, and are of excellent eating quality. The new variety ripens midseason, before any other muscat flavored, seedless grape cultivars grown in California.

20 The new variety was first hybridized by David W. Cain and Michael J. Striem in Wasco, Kern County, California, the variety being originated by controlled hybridization and subsequent ovule culture of seed traces and abortive embryo. The seed parent is the variety 'Black Monukka' (unpatented), and the pollen parent is 'Sugrafive' (U.S. Plant Patent No. 5,151). The parent varieties were first crossed in
25 May 1988, with the date of first sowing being August 1988 and the date of first flowering being May, 1991. The new 'Sugratwentyfour' variety was first asexually propagated by David W. Cain in January, 1992, in Wasco, Kern County, using hardwood cuttings.

30 'Sugratwentyfour' is distinguished from other commonly grown white seedless grapevine varieties such as 'Sugraeighteen', 'Sugraone', 'Perlette', and 'Thompson Seedless'.

The new grapevine variety 'Sugratwentyfour' resembles the 'Sugraeighteen' grapevine variety by its distinct muscat flavor, berry shape and color. The new grapevine variety differs from the 'Sugraeighteen' variety by having a substantially earlier ripening period – grapes of 'Sugratwentyfour' ripen about the same time as 'Sugraone' and 'Thompson Seedless' varieties. Additionally, the vine of the new variety is more vigorous and less productive (cane pruned) than 'Sugraeighteen' (spur pruned).

The new grapevine variety 'Sugratwentyfour' differs from the 'Sugraone', 'Perlette' and 'Thompson Seedless' and other commonly grown white seedless grapevine varieties by possessing a distinct and moderately mild muscat flavor somewhat similar to that of the seeded Italia grapevine varieties. It also differs from the above listed varieties by having much more nearly round berries than 'Sugraone' or 'Thompson Seedless'.

The new 'Sugratwentyfour' variety has been shown to maintain its distinguishing characteristics through successive asexual propagations by, for example, cuttings.

BRIEF DESCRIPTION OF THE FIGURE

The accompanying photographic illustration in Fig. 1 illustrates in full color a typical cluster of berries, a young shoot, and a mature leaf blade of the new grapevine. The colors are as nearly true as is reasonably possible in a color representation of this type.

DETAILED BOTANICAL DESCRIPTION OF THE INVENTION

Throughout this specification, color names beginning with a small letter signify that the name of that color, as used in common speech, is aptly descriptive. Color names beginning with a capital letter designate values based upon the R.H.S. Colour Chart, published by The Royal Horticultural Society, London, England.

Many of the description values in this specification are based on and conform to those set forth by the International Board for Plant Genetic Resources Institute Grape Descriptors (*Vitis* spp.) of 1983 and/or 1997 which was developed in collaboration with

the Office International de la Vigne et du Vin (OIV) and the International Union for the Protection of New Varieties of Plants (UPOV).

5 The descriptive matter which follows pertains to 'Sugratwentyfour' plants grown in the vicinity of Wasco, Kern County, California, during 2002, and is believed to apply to plants of the variety grown under similar conditions of soil and climate elsewhere.

VINE

General:

10 *Size:* medium.
Vigor: medium.
Density of foliage: medium.
Productivity: medium productive.
Root stock: own roots.

Trunk:

15 *Shape:* slender.
Straps: long.
Surface texture: medium.
Inner bark color: about Greyed-orange 177B.

20 SHOOTS

Young shoot:

Form of tip: half-open.
Distribution of anthocyanin coloration of tip: absent.
Intensity of anthocyanin coloration of tip: absent.
25 *Density of prostrate hairs on tip:* very sparse.
Density of erect hairs on tip: absent.

Woody Shoot (Mature Canes):

Shape: medium.
Internode length: long - approximately 35.7 mm.
30 *Width at node:* approximately 14.83 mm.
Cross section: circular.

Surface: ribbed.

Main color: light brown.

Lenticels: absent.

Density of erect hairs on nodes: none or very sparse.

5 *Density of erect hairs on internodes:* none or very sparse.

Growth of axillary shoots: weak; approximately 17.9 cm.

Flowering shoot:

Vigor during flowering: strong.

Attitude during flowering on shoots which are not tied: semi-erect.

10 *Color of dorsal side of internodes:* about Yellow-green 144A.

Color of ventral side of internodes: about Yellow-green 144A.

Color of dorsal side of nodes: about Yellow-green 144A.

Color of ventral side of nodes: about Yellow-green 144A.

Density of erect hairs on nodes: absent.

15 *Erect hairs on internode:* absent.

Density of prostrate hairs on nodes: absent.

Density of prostrate hairs on internodes: absent.

Anthocyanin coloration of buds: absent.

Tendrils:

20 *Distribution on the shoot at full flowering:* discontinuous.

Thickness: thick.

Color: about Yellow-green 151A.

Form: bifurcated.

Number of consecutive tendrils: up to two.

25 *Length of tendril:* medium; approximately 19 cm.

LEAVES

Young Leaves:

Color of upper surface of first 4 distal unfolded leaves: green.

30 *Average intensity of anthocyanin coloration of six distal leaves prior to flowering:* absent.

Density of prostrate hairs between veins at lower surface of 4th distal unfolded leaf: absent.

Density of erect hairs between veins at lower surface of 4th distal unfolded leaf: absent.

5 *Density of prostrate hairs on veins at lower surface of 4th distal unfolded leaf: absent.*

Density of erect hairs on veins at lower surface of 4th distal unfolded leaf: absent.

Mature leaves:

10 *Average length: approximately 103 mm.*

Average width: approximately 164 mm.

Size of blade: medium.

Shape of blade: pentagonal.

Number of lobes: three.

15 *Anthocyanin coloration of main veins on the upper side of the blade: absent.*

Mature leaf profile: flat.

Blistering surface of blade upper surface: absent.

Leaf blade tip: curved downwardly.

Undulation of margin: slight.

20 *Apex: cuspidate.*

Thickness: medium.

Undulation of blade between main and lateral veins: absent.

Shape of teeth: mixture of both sides straight and both sides convex.

Length of teeth: short.

25 *Ratio length/width of teeth: small.*

General shape of petiole sinus: half-open.

Tooth at petiole sinus: absent.

Petiole sinus limited by veins: absent.

Shape of upper lateral sinus: open.

30 *Depth of upper lateral sinus: shallow.*

Density of prostrate hairs between veins on lower surface of blade: absent.

Density of erect hairs between veins on lower surface of blade: absent.

Density of prostrate hairs on main veins on lower surface of blade: none or very sparse.

5 *Density of erect hairs on main veins on lower surface of blade:* none or very sparse.

Density of prostrate hairs on main veins on upper surface of blade: absent.

Autumn coloration of leaves: about Yellow-green 153B.

Upper Surface:

Color: about Yellow-green 147A.

10 *Surface texture:* smooth .

Surface appearance: semi-glossy.

Goffering of blade: absent.

Lower Surface:

Color: about Yellow-green 146A.

15 *Anthocyanin coloration of main veins on lower leaf surface:* absent.

Glossiness: weak.

Pubescence: absent.

Surface texture: wrinkled.

Surface appearance: dull.

20 **Petiole:**

Length of petiole: medium; approximately 9.5 cm.

Length of petiole compared to middle vein: slightly shorter.

Density of prostrate hairs on petiole: absent.

Density of erect hairs on petiole: absent.

25 *Shape of base of petiole sinus:* U-shaped.

Buds:

Shape: slightly pointed.

Size: medium; approximately 4 by 6 mm.

Position: slightly held out.

30 *Cane bud fruitfulness:* basal most fruitful.

Time of bud burst: early – March 18, 2002.

FLOWERS

General:

Flower sex: hermaphrodite.

5 *Length of first inflorescence:* medium.

Position of first flowering node: second node.

Number of inflorescences per shoot: approximately 1.1 to 2.

Date of full bloom: May 11, 2002.

10 *Time of bloom:* early (as compared to similar varieties growing in the same region.)

Size (diameter of fully open flower): medium.

FRUIT

General:

15 *Ripening period:* medium – approximately 0 days ahead of Thompson Seedless
variety.

Use: fresh market.

Keeping quality: medium.

Resistance: *Insects:* medium.

20 *Diseases: medium.*

Shipping quality: medium.

Date of first harvest: July 15, 2002.

Solids-sugar: medium - approximately 18%.

Refractometer test: approximately 18.0.

25 *Acid*: medium – approximately 5.04 g/liter.

Juice pH: approximately 3.9.

Cluster:

Bunch size (peduncle excluded): medium.

Bunch length (peduncle excluded): intermediate – approximately 19.4 cm.

30 *Bunch width:* approximately 16.1 cm.

Bunch weight: low – approximately 561 g.

Bunch density: loose.

Number of berries: approximately 152.

Form: cylindrical.

Peduncle:

5 *Length of peduncle:* short – approximately 4.07 cm.

Lignification of peduncle: weak.

Color: about Yellow-green 146B.

Berry:

Size: medium.

10 *Uniformity of size:* variable.

Berry weight: low – approximately 3.68g.

Shape: round.

Presence of seeds: rudimentary - approximately 0.6 mg.

Cross section: circular.

15 *Dimensions:* approximately 198 mm longitudinal axis by approximately 168 mm
 horizontal axis.

Skin color (without bloom): about Greyed-yellow 160A.

Coloration of flesh: transparent greenish.

Juiciness of flesh: slightly juicy.

20 *Berry firmness:* medium.

Particular flavor: muscat.

Bloom (cuticular wax): very weak.

Pedice length: approximately 7.80 mm.

Berry separation from pedicel: easy.

25 *Visibility of hilum:* slightly clear.

Skin:

Thickness: medium.

Texture: medium.

Reticulation: absent.

30 *Roughness:* absent.

Tenacity: free.